

q-Deformed Superconformal Algebra on Quantum Superspace

Tatsuo Kobayashi* and Tsuneo Uematsu†

Department of Fundamental Sciences, FIHS,
Kyoto University, Kyoto 606, Japan

Abstract

A quantum deformation of 4-dimensional superconformal algebra realized on quantum superspace is investigated. We study the differential calculus and the action of the quantum generators corresponding to $sl_q(1|4)$ which act on the quantum superspace. We derive deformed $su(1|2, 2)$ algebras from the deformed $sl(1|4)$ algebra. Through a contraction procedure we obtain a deformed super-Poincaré algebra.

* Fellow of the Japan Society for the Promotion of Science. Work partially supported by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture (# 030083)

†Work partially supported by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture (# 04245221)

Recently there has been a great deal of interest in quantum groups and quantum algebras [1, 2, 3, 4] in various fields of theoretical physics and mathematics, such as conformal field theory, integral models, statistical mechanics, knot theory and so on [5, 6, 7, 8].

Quantum groups can be realized on a quantum space or a quantum (hyper-)plane in which coordinates are non-commuting [7, 9]. The differential calculus on the non-commutative space [10] has been developed by Wess and Zumino [11] and by other people [12, 13, 14], especially for $GL_q(n)$ [12, 13, 14], as well as for $SO_q(n)$ [15].

So far most of the quantum deformations have been applied to the Lie groups or algebras corresponding to the internal symmetries. Now it would be interesting to explore quantum deformation of space-time symmetries, which might be relevant at very short distances e.g. at the Planck length implied by unification theories including gravity. In the last few years, there have been pioneering works in extending quantum-group ideas to space-time symmetries including Lorentz, Poincaré and conformal algebras [16–23].

In our previous paper [24], we studied a quantum deformation of 4-dimensional conformal algebras based on the quantum space realizing $sl_q(4)$. Through a contraction procedure we also derived a quantum deformation of Poincaré algebra. In the present paper, we shall investigate a q-deformed superconformal algebra in 4-dimensions in the framework of the quantum superspace which

realizes q-deformed superalgebra, $sl_q(1|4)$. By taking a contraction limit we obtain a deformed super-Poincaré algebra.

Following [11, 24] we now consider actions of deformed $sl(1|4)$ generators on a quantum superspace, whose coordinates Z^I and derivatives ∂_I consist of bosonic elements x^0, ∂_0 and fermionic ones $\theta^\alpha, \partial_\alpha$ ($\alpha = 1 \sim 4$). In Ref.[24], the deformed $sl(4)$ algebra has been represented on a 4-dimensional quantum bosonic space. In this paper the same deformed algebra shall be obtained on the fermionic space to supersymmetrize the algebra.

Differential calculus on general quantum superspaces with multi-parameters has been studied in [25]. Here we set up the following commutation relations of the

coordinates and the derivaties with one-parameter,[‡]

$$\begin{aligned}
Z^I Z^J &= \frac{(-1)^{\hat{I}\hat{J}}}{q} Z^J Z^I, \quad \partial_I \partial_J = (-1)^{\hat{I}\hat{J}} q \partial_J \partial_I, \quad (I < J), \\
\partial_I Z^J &= \frac{(-1)^{\hat{I}\hat{J}}}{q} Z^J \partial_I, \quad (I \neq J), \\
\partial_I Z^I &= 1 - (-q)^{2\hat{I}-2} Z^I \partial_I + (q^2 - 1) \sum_{I < J} Z^J \partial_J,
\end{aligned} \tag{1}$$

where $\hat{I}$ denotes the grassmann parity, i.e., $\hat{I} = 0$ for $I = 0$ and $\hat{I} = 1$ for $I \neq 0$. These commutation relations are written in terms of the following $\hat{R}$ -matrix,

$$\hat{R}^{IJ}_{KL} = \delta_L^I \delta_K^J \{ \delta^{IJ} ((-q)^{2\hat{I}} - (-1)^{\hat{I}\hat{J}} q) + (-1)^{\hat{I}\hat{J}} q \} + \delta_K^I \delta_L^J \Theta^{JI} (1 - q^2), \tag{2}$$

where $\Theta^{IJ} = 1$ for $I > J$ otherwise it vanishes. For example, we can use the $\hat{R}$ -matrix to write the commutation relations between Z^I and ∂_K as

$$\partial_K Z^I = \delta_K^I + q^{-2} \hat{R}^{IJ}_{KL} Z^L \partial_J. \tag{3}$$

Ordering of the bosonic and fermionic coordinates is non-trivial due to Θ . We study here ordering where θ^α follows x^0 , i.e., $\Theta^{0\alpha} = 0$ and $\Theta^{\alpha 0} = 1$. We could discuss the other ordering.

Now we consider actions of the deformed $sl(1|4)$ generators on the above quantum space. First of all, we discuss actions of T^I_J ($J = I + 1, I - 1$) which correspond to $Z^I \partial_J$ in the “classical” limit ($q \rightarrow 1$). Following [11], we choose the following basis of the algebra,

$$[T^I_J, Z^K]_{a(I,J,K)} = \delta_J^K Z^I, \quad [T^I_J, \partial_K]_{b(I,J,K)} = c(I, J) \delta_K^I \partial_J, \tag{4}$$

where $[A, B]_a \equiv AB - aBA$ and it is called a q -commutator hereafter. Similarly, a q -anticommutator is defined as $\{A, B\}_a \equiv AB + aBA$. We investigate consistency between (1) and (4), so as to obtain the following solution,

$$a(I, J, K) = b(I, J, K)^{-1} = -c(I, I + 1)^{-1} = q^{2\hat{I}-1}, \quad (K = \min(I, J))$$

$$a(I, J, K)^{-1} = b(I, J, K) = c(I + 1, I) = (-1)^{\hat{I}+1} q, \quad (K = \max(I, J))$$

[‡]Matrices which transform covariantly these commutation relations have a superdeterminant as a center (see in detail [25]).

$$a(I, J, K) = b(I, J, K) = (-1)^{\hat{K}(\hat{I}+\hat{J})}, \quad (K \neq I, J), \quad (5)$$

where $J = I \pm 1$ and $\min(I, J)$ ($\max(I, J)$) denotes the smaller (larger) number in the parenthesis.

Next we define Cartan elements H_I ($I = 0 \sim 3$) using q -commutators of T_{I+1}^I and T_I^{I+1} . We choose H_I as follows,

$$H_0 = \{T_0^1, T_1^0\}, \quad H_\alpha = q^{-1}[T_\alpha^{\alpha+1}, T_{\alpha+1}^\alpha]_{q^2}, \quad (6)$$

for the purpose that terms depending on T_K^I disappear in the actions of H_I . Note that in the classical limit $H_I = T_{I+1}^{I+1} + (-1)^{\hat{I}} T_I^I = Z^{I+1} \partial_{I+1} + (-1)^{\hat{I}} Z^I \partial_I$. It is remarkable that the definition of H_0 is not deformed. The Cartan elements act on the quantum space as follows,

$$\begin{aligned} [H_0, Z^I]_{1/q^2} &= Z^I, \quad [H_0, \partial_I]_{q^2} = -q^2 \partial_I, \quad (I = 0, 1), \\ [H_\alpha, \theta^\alpha]_{q^2} &= -q\theta, \quad [H_\alpha, \theta^{\alpha+1}]_{1/q^2} = q^{-1} \theta^{\alpha+1}, \\ [H_\alpha, \partial_\alpha]_{1/q^2} &= q^{-1} \partial_\alpha, \quad [H_\alpha, \partial_{\alpha+1}]_{q^2} = -q \partial_{\alpha+1}, \end{aligned} \quad (7)$$

and H_I commute with the other coordinates and the other derivatives.

The other elements of the deformed $sl(1|4)$ can be defined in terms of

q -commutators of T_{I+1}^I and T_J^{J+1} so that their commutation relations close. The requirement of the algebraic closure is satisfied if we choose the other elements as follows,

$$T_K^I = [T_J^I, T_K^J]_q, \quad (I < J < K \text{ or } I > J > K). \quad (8)$$

The general non-Cartan generators act on the quantum space as follows,

$$\begin{aligned} T_K^I Z^K &= -(-1)^{\hat{I}} q^{-1} Z^K T_K^I + Z^I + (-1)^{\hat{I}} (q - q^{-1}) \sum Z^J T_J^I, \\ T_I^K Z^I &= q^{2\hat{I}-1} Z^I T_I^K + Z^K, \quad T_I^K Z^J = -(-1)^{\hat{I}} (Z^J T_I^K + (q - q^{-1}) Z^K T_I^J) \\ [T_\alpha^0, x^0]_{1/q} &= \{T_\alpha^\alpha, \theta^\alpha\}_{1/q} = [T_\beta^\alpha, \theta^\alpha]_q = [T_\alpha^\beta, \theta^\beta]_{1/q} = 0, \\ T_K^I \partial_I &= q^{1-2\hat{I}} (\partial_I T_K^I - \partial_K), \quad T_K^I \partial_J = -(-1)^{\hat{I}} \partial_J T_K^I + (q - q^{-1}) \partial_K T_J^I, \\ T_I^K \partial_K &= -(-1)^{\hat{I}} \{ \partial_K T_I^K + q^{2(K-I)-1} \partial_I - (q - q^{-1}) \sum q^{2(K-J)} \partial_J T_I^J \}, \\ \{T_\alpha^0, \partial_\alpha\}_q &= [T_\alpha^\alpha, \partial_0]_q = [T_\beta^\alpha, \partial_\beta]_q = [T_\alpha^\beta, \partial_\alpha]_{1/q} = 0, \end{aligned} \quad (9)$$

where $I < J < K$ and $\alpha < \beta$, and the non-Cartan generators and the other elements of the quantum space satisfy the “classical” algebra, i.e., they commute or anti-commute

each other, depending upon their grassmann parity. Using (7) and (9), we can derive commutation relations between the deformed $sl(1|4)$ generators. Some commutation relations among the whole deformed $sl(1|4)$ algebra are shown in the following;

$$\begin{aligned}
[T_{I+1}^I, T_{I+1}^{I+2}]_q &= [T_{I+2}^{I+1}, T_I^{I+1}]_q = [T_J^I, T_L^K] = 0, \quad (I, J < K, L), \\
[H_0, T_\alpha^1]_{1/q^2} &= T_\alpha^1, \quad [H_0, T_1^\alpha]_{q^2} = -q^2 T_1^\alpha, \\
[H_\alpha, T_\alpha^0]_{1/q^2} &= q^{-1} T_\alpha^1, \quad [H_\alpha, T_0^\alpha]_{q^2} = -q T_0^\alpha, \\
[H_\alpha, T_{\alpha+1}^\alpha]_{q^4} &= -q^2(q + q^{-1}) T_{\alpha+1}^\alpha, \quad [H_\alpha, T_{\alpha+1}^{\alpha+1}]_{1/q^4} = q^2(q + q^{-1}) T_{\alpha+1}^{\alpha+1}, \\
[H_\alpha, T_{\beta+1}^\beta]_{1/q^2} &= q^{-1} T_{\beta+1}^\beta, \quad [H_\alpha, T_\beta^{\beta+1}]_{q^2} = -q T_\beta^{\beta+1}, \quad (\alpha = \beta \pm 1), \\
[H_0, T_0^1] &= [H_0, T_1^0] = [H_I, T_K^J] = 0, \quad (J, K \neq I, I+1),
\end{aligned} \tag{10}$$

and the other commutation relations of the deformed $sl(1|4)$ algebra can be derived from the above commutation relations. It is clear that the Cartan elements commute with each other and T_α^0 and T_0^α are nilpotent. In this algebra, the generators H_α and T_β^α construct $sl_q(4)$ algebra as a closed subalgebra of the deformed $sl(1|4)$. It implies that from the algebra we can derive a deformed superconformal algebra which includes the deformed conformal algebra as a closed subalgebra. The whole commutation relations of the deformed $sl(4)$ algebra have been shown explicitly in [24] and the whole commutation relations of the deformed $sl(1|4)$ algebra will be explicitly shown elsewhere.

From the present result, we can easily derive the deformed superconformal algebra in 2-dimensions, $sl_q(1|2)$, by restricting ourselves to the generators H_0, H_1, T^I_J ($I, J = 0, 1, 2$).

We need another space conjugate to Z^I in order to define the deformed superconformal algebra $su(1|2, 2)$ from the deformed $sl(1|4)$ algebra. Here we introduce the conjugate space $\bar{Z}_I = (\bar{x}_0, \bar{\theta}_\alpha)$. Using the $\hat{R}$ -matrix, we set up commutation relations of $\bar{Z}_I$ as follows;

$$\bar{Z}_L \bar{Z}_K = \hat{R}_{KL}^{IJ} \bar{Z}_J \bar{Z}_I, \quad \bar{Z}_k Z^I = \hat{R}_{KL}^{IJ} Z^L \bar{Z}_J. \tag{11}$$

These commutation relations have a center $C = x^0 \bar{x}_0 + \sum_\alpha \theta^\alpha \bar{\theta}_\alpha$. Further, we assume that the deformed $sl(1|4)$ generators have the same actions on $\bar{Z}_I$ as those on ∂_I . For example, the generator T_K^I ($I < K$) acts on $\bar{Z}_I$ as follows,

$$T_K^I \bar{Z}_I = q^{1-2\hat{I}} (\bar{Z}_I T_K^I - \bar{Z}_K). \tag{12}$$

Now we consider conjugation between Z^I and $\bar{Z}_I$. We assume that Z^I and $\bar{Z}_I$ are related as

$$\bar{x}_0 = K_0(x^0)^*, \quad \bar{\theta}_\alpha = K_\alpha \eta^\alpha (\theta^\alpha)^*, \quad (13)$$

where $*$ denotes complex conjugate and K_α are coefficients which become identity in the classical limit and η^α is a $su(2, 2)$ metric, i.e., $\eta^\alpha = (1, 1, -1, -1)$. The conjugation (13) should be consistent with the commutation relations, (1) and (11). We can find two types of the consistent conjugations depending upon a value of the deformation parameter q . At first we discuss the case where q is real. In this case, the conjugation is allowed if $K_I = 1$ and the conjugation includes reversing of order, i.e., $\overline{ab} = \bar{b}\bar{a}$. Then we take the conjugation of (12) and compare it with (9), so that we obtain conjugation relations of the generators as follows,

$$\overline{T^I_J} = \eta^I \eta^J T^J_I. \quad (14)$$

Similarly, we have

$$\overline{H_I} = H_I. \quad (15)$$

Next we discuss the case with $|q| = 1$, where the conjugation does not include reversing of order, i.e., $\overline{ab} = \bar{a}\bar{b}$. This conjugation is consistent with (1) and (11) if $K_\alpha = q^\alpha$ and $K_0 = iq$. Further we follow the procedure similar to (14) to obtain the following conjugation relations;

$$\overline{T^I_J} = -q^{I-J+1} \eta^I \eta^J T^J_I, \quad \overline{H_I} = -q^{2I-2} H_I, \quad (I < J) \quad (16)$$

where $\eta^0 = i/q$.

We now obtain the quantum deformation of 4-dimensinal superconformal algebra by assigning the $su_q(1|2, 2)$ elements to the physical superconformal generators. Here we shall consider the case where the deformation parameter q is real. The similar analysis can be performed for the case of $|q| = 1$.

We first consider the generators of the conformal sector. The $SO(4, 2)$ generators M_{mn} ($m, n = 0, 1, 2, 3, 4, 5$) with the metric $g_{mn} = \text{diag}(1, -1, -1, -1, -1, 1)$ satisfy

$$[M_{mn}, M_{kl}] = -ig_{mk} M_{nl} + ig_{nk} M_{ml} + ig_{ml} M_{nk} - ig_{nl} M_{mk}. \quad (17)$$

This can be realized by the $su(1|2, 2)$ elements as $M_{mn} = T^\alpha_\beta (\sigma_{mn})_\alpha^\beta$ (See e.g. ref.[26]) where we take $(\mu, \nu = 0, 1, 2, 3; \alpha, \beta = 1, \dots, 4)$

$$\begin{aligned} \sigma_{\mu\nu} &= \frac{i}{4} [\gamma_\mu, \gamma_\nu], & \sigma_{4\mu} &= \frac{i}{2} \gamma_\mu, \\ \sigma_{5\mu} &= \frac{i}{2} \gamma_5 \gamma_\mu, & \sigma_{45} &= \frac{i}{2} \gamma_5. \end{aligned} \quad (18)$$

The conjugation of T^α_β (14) and (15) for q real, leads to the reality of M_{mn} :

$$\overline{M_{mn}} = M_{mn} \quad (19)$$

with the suitable representation for gamma-matrices γ_μ ($\mu = 0, 1, 2, 3$) which satisfy

$$\eta^\alpha \eta^\beta (\sigma^*_{mn})_\alpha{}^\beta = (\sigma_{mn})_\beta{}^\alpha. \quad (20)$$

Now we regard the 4-th and 5-th dimensions as the compact space. We construct the generators of classical $4D$ conformal group out of these rotational generators of $SO(4, 2)$. First, the Lorentz rotation operators $M_{\mu\nu}$ ($\mu, \nu = 0, 1, 2, 3$) is given as $M_{\mu\nu} = M_{m=\mu, n=\nu}$ with

$$\begin{aligned} M_{12} &= \frac{-1}{2}(H_1 + H_3), & M_{23} &= \frac{-1}{2}(T^1_2 + T^2_1 + T^3_4 + T^4_3), \\ M_{31} &= \frac{i}{2}(T^1_2 - T^2_1 + T^3_4 - T^4_3), & M_{01} &= \frac{-1}{2}(T^1_4 - T^4_1 + T^2_3 - T^3_2) \\ M_{02} &= \frac{i}{2}(T^1_4 + T^4_1 - T^2_3 - T^3_2), & M_{03} &= \frac{-1}{2}(-T^1_3 + T^3_1 + T^2_4 - T^4_2) \end{aligned} \quad (21)$$

while the translation operators P_μ , conformal boost operators K_μ and dilatation operator D are given as $P_\mu = M_{4\mu} + M_{5\mu}$, $K_\mu = M_{5\mu} - M_{4\mu}$, $D = M_{45}$ with

$$\begin{aligned} M_{40} &= \frac{-i}{2}(T^1_3 + T^3_1 + T^2_4 + T^4_2), & M_{41} &= \frac{-1}{2}(T^1_2 + T^2_1 - T^3_4 - T^4_3) \\ M_{42} &= \frac{i}{2}(T^1_2 - T^2_1 - T^3_4 + T^4_3), & M_{43} &= \frac{-1}{2}(H_1 - H_3) \\ M_{50} &= \frac{-1}{2}(H_1 + 2H_2 + H_3), & M_{51} &= \frac{i}{2}(T^2_3 + T^3_2 + T^1_4 + T^4_1) \\ M_{52} &= \frac{-1}{2}(T^2_3 - T^3_2 - T^1_4 + T^4_1), & M_{53} &= \frac{i}{2}(-T^1_3 - T^3_1 + T^2_4 + T^4_2) \\ M_{45} &= \frac{-1}{2}(T^1_3 - T^3_1 + T^2_4 - T^4_2), \end{aligned} \quad (22)$$

We assign the generators T^0_β and T^β_0 to the supercharges Q_α , $\overline{Q}_{\dot{\alpha}}$, S^α , $\overline{S}^{\dot{\alpha}}$ as follows:

$$\begin{aligned} Q_1 &= \sqrt{2}(T^0_1 - iT^0_3), & \overline{Q}_1 &= \sqrt{2}(T^1_0 - iT^3_0) \\ Q_2 &= \sqrt{2}(-T^0_2 + iT^0_4), & \overline{Q}_2 &= \sqrt{2}(-T^2_0 + iT^4_0) \\ \overline{S}^1 &= \sqrt{2}(T^0_1 + iT^0_3), & S^1 &= \sqrt{2}(T^1_0 + iT^3_0) \\ \overline{S}^2 &= \sqrt{2}(-T^0_2 - iT^0_4), & S^2 &= \sqrt{2}(-T^2_0 - iT^4_0) \end{aligned} \quad (23)$$

Note that here we have the conjugation (14). In 4-dimensional $N=1$ superconformal algebra, in addition to $M_{\mu\nu}$, P_μ , K_μ and D we have another bosonic generator, the $U(1)$

charge, which is defined as

$$A = -\frac{1}{4}(4H_0 + 3H_1 + 2H_2 + H_3) \quad (24)$$

Now we take the same assignment of the generators for the q -deformed case. The q -deformed superconformal algebra can be read off from the commutation relations among the $sl_q(1|4)$ generators.

The classical anticommutation relation of Q_α and $\bar{Q}_{\dot{\beta}}$ given by

$$\{Q_\alpha, \bar{Q}_{\dot{\beta}}\} = 2(\sigma^\mu)_{\alpha\dot{\beta}} P_\mu \quad (25)$$

where $(\sigma^\mu) = (\mathbf{1}, \sigma^i)$ and $(\bar{\sigma}^\mu) = (\mathbf{1}, -\sigma^i)$, is deformed as, for example for $\alpha = \dot{\beta} = 1$,

$$\begin{aligned} \{Q_1, \bar{Q}_1\} = & 2P_+ + 2[(1 - q^4)H_0 + (1 - q^3)H_1 + (1 - q)H_2 + i(1 - q)T^1_3 \\ & + i(1 - q^3)T^3_1 + i\lambda T^1_3 H_0 - i(1 - q)T^1_0 T^0_3 + iq^2 \lambda T^3_1 H_0 + q^3 \lambda T^0_1 T^1_0 \\ & + q\lambda T^0_2 T^2_0 + q\lambda H_0 H_1 + \lambda H_2 H_0 + q^3 \lambda T^2_3 T^3_2 + q^3 \lambda T^1_3 T^3_1 + q\lambda H_1 H_2 \\ & + q^4 \lambda^2 T^1_2 T^2_1 - q^3 \lambda^2 T^2_3 T^3_2 H_1 - q^2 \lambda^2 T^1_3 T^3_1 H_0 - q^2 \lambda^2 T^2_3 T^3_2 H_0 \\ & - \lambda^2 H_1 H_2 H_0 + q^2 \lambda^3 T^2_3 T^3_2 H_1 H_0 - q^3 \lambda^3 T^1_2 T^2_1 H_0] \end{aligned} \quad (26)$$

where $P_\pm = P_0 \pm P_3$ and $\lambda = q - q^{-1}$. The classical relation $\{\bar{S}^{\dot{\alpha}}, S^\beta\} = 2(\bar{\sigma}^\mu K_\mu)^{\dot{\alpha}\beta}$ is similarly deformed. While another classical anticommutation relation

$$\{Q_\alpha, S^\beta\} = (\sigma^{\mu\nu})_\alpha{}^\beta M_{\mu\nu} + 2i\delta_\alpha{}^\beta D - 4\delta_\alpha{}^\beta A \quad (27)$$

receives q -deformation, e.g. for $\alpha = \beta = 1$, as given by

$$\begin{aligned} \{Q_1, S^1\} = & (\sigma^{\mu\nu})_1{}^1 M_{\mu\nu} + 2iD - 4A + 2[(q^4 - 1)H_0 + (q^3 - 1)H_1 + (q - 1)H_2 \\ & + i(1 - q)T^1_3 - i(1 - q^3)T^3_1 + i\lambda T^1_3 H_0 - i(1 - q)T^1_0 T^0_3 - iq^2 \lambda T^3_1 H_0 \\ & - q^3 \lambda T^0_1 T^1_0 - q\lambda T^0_2 T^2_0 - q\lambda H_0 H_1 - \lambda H_2 H_0 - q^3 \lambda T^2_3 T^3_2 - q^3 \lambda T^1_3 T^3_1 \\ & - q\lambda H_1 H_2 - q^4 \lambda^2 T^1_2 T^2_1 + q^3 \lambda^2 T^2_3 T^3_2 H_1 + q^2 \lambda^2 T^1_3 T^3_1 H_0 + q^2 \lambda^2 T^2_3 T^3_2 H_0 \\ & + \lambda^2 H_1 H_2 H_0 - q^2 \lambda^3 T^2_3 T^3_2 H_1 H_0 + q^3 \lambda^3 T^1_2 T^2_1 H_0] \end{aligned} \quad (28)$$

where $\sigma^{\mu\nu} = \frac{i}{2}(\sigma^\mu \bar{\sigma}^\nu - \sigma^\nu \bar{\sigma}^\mu)$ and $\bar{\sigma}^{\mu\nu} = \frac{i}{2}(\bar{\sigma}^\mu \sigma^\nu - \bar{\sigma}^\nu \sigma^\mu)$.

The anti-commutator $\{Q_\alpha, Q_\beta\}$ is also deformed. We find

$$\{Q_1, Q_2\}_q = 2(1 - q^2)T^0_3 T^0_2, \quad Q_1^2 = 2i(q - 1)T^0_3 T^0_1, \quad Q_2^2 = 2i(q - 1)T^0_4 T^0_2, \quad (29)$$

where we note that Q_α is no more nilpotent. We further observe that $Q_\alpha^3 = 0$, which has a common feature with parafermion statistics. It would be very interesting to investigate the representation theory of this q -deformed algebra.

Since super-Poincaré algebra is not a simple Lie superalgebra, we cannot apply the standard prescription of quantum deformation. As in the case of Poincaré algebra, we need some contraction procedure. Here we shall adopt the contraction method using rescaling of the translation as well as supersymmetry generators and subsequent limiting procedure [27].

We first make the following rescaling of the generators

$$\begin{aligned}\hat{M}_{\mu\nu} &= M_{\mu\nu}, & \hat{P}_\mu &= \frac{1}{R}P_\mu, & \hat{K}_\mu &= K_\mu, & \hat{D} &= D, & \hat{A} &= A, \\ \hat{Q}_\alpha &= \frac{1}{\sqrt{R}}Q_\alpha, & \hat{\bar{Q}}_{\dot{\alpha}} &= \frac{1}{\sqrt{R}}\bar{Q}_{\dot{\alpha}}, & \hat{S}^\alpha &= S^\alpha, & \hat{\bar{S}}^{\dot{\alpha}} &= \bar{S}^{\dot{\alpha}}.\end{aligned}\tag{30}$$

Let us substitute the above rescaled generators into the q -deformed superconformal algebra and then take the contraction limit:

$$R \rightarrow \infty, \quad q \rightarrow 1,\tag{31}$$

with $\kappa = (q - q^{-1})R^2$ kept fixed. The κ is a new deformation parameter similar to those introduced in refs. [21, 22, 23, 27, 24].

We then get a quantum deformation of super-Poincaré algebra. In the following we omit the hat again. In the sector of the bosonic generators, only the commutation relations among the Lorentz generators, $[M_{\mu\nu}, M_{\lambda\sigma}]$, receive the deformation which was discussed in the previous paper [24]. The other commutation relations, e.g., $[M_{\mu\nu}, P_\sigma]$, $[P_\mu, P_\nu]$ and $\{Q_\alpha, \bar{Q}_{\dot{\beta}}\}$ etc. remain classical.

In this paper we have investigated the q -deformed $4D$ superconformal and super-Poincaré algebras in the framework of the quantum space for $sl_q(1|4)$. We studied the differential calculus on the non-commutative quantum space for $sl_q(1|4)$ as well as the action of the generators on this quantum space. Through the charge conjugations, we obtain two types of the deformed $su(1|2, 2)$ algebras from the deformed $sl(1|4)$.

We presented the q -deformed superconformal algebra with a suitable assignment of the $4D$ superconformal generators. The algebra includes the deformed conformal algebra as a closed subalgebra. A quantum super-Poincaré algebra with a deformation parameter κ was obtained by rescaling of the translation and supercharge operators and subsequent contraction procedure.

In the present paper we only discussed the 4-dimensional case corresponding to $sl_q(1|4)$. As we mentioned before, it is quite easy to derive the 2-dimensional case, $sl_q(1|2)$, which is a deformation of the subalgebra of super Virasoro algebra. As a future problem, it would be very much interesting to extend the present analysis to the extended superconformal algebras.

Acknowledgement

The authors would like to thank P. P. Kulish, R. Sasaki and C. Schwiebert for valuable discussions.

References

- [1] V. G. Drinfeld, Sov. Math. Dokl. **32**(1985)254.
- [2] M. Jimbo, Lett. Math. Phys. **10**(1985)63.
- [3] L. D. Faddeev, N. Yu. Reshetikhin and L. A. Takhtajan, Algebra and Analysis, **1**(1987)178.
- [4] P. P. Kulish and N. Yu. Reshetikhin, J. Sov. Math. **23**(1983)2435.
- [5] H. -D. Doebner and J. -D. Hennig eds., *Quantum Groups*, Lecture Notes in Physics **370**(Springer Verlag, 1990).
- [6] T. Curtright, D. Fairlie and C. Zachos eds., *Quantum Groups*, Proceedings of the Argonne Workshop (World Scientific, 1990).
- [7] L. A. Takhtajan, Adv. Studies Pure Math. **19**(1989)435; Lecture Notes in Physics **370**(1989)3.
- [8] P. P. Kulish ed., *Quantum Groups*, Lecture Notes in Mathematics **1510**(Springer Verlag, 1992).
- [9] Yu. I. Manin, Commun. Math. Phys. **123**(1989)163.
- [10] S. L. Woronowicz, Commun. Math. Phys. **111**(1987)613; **122**(1989)125.
Publ. RIMS, Kyoto Univ. **23**(1987)117.
- [11] J. Wess and B. Zumino, Nucl. Phys. B (Proc. Suppl.) **18B**(1990)302.
- [12] A. Schirmacher, J. Wess and B. Zumino, Z. Phys. C – Particles and Fields **49**(1991)317.
- [13] A. Schirmacher, Z. Phys. C – Particles and Fields **50**(1991)321.
- [14] A. Schirmacher, J. Phys. A: Math. Gen. **24**(1991)L1249.
- [15] U. Carow-Watamura, M. Schlieker and S. Watamura, Z. Phys. C – Particles and Fields **49**(1991) 439.
- [16] P. Podleś and S. L. Woronowicz, Commun. Math. Phys. **130**(1990)381.

- [17] U. Carow-Watamura, M. Schlieker, M. Scholl and S. Watamura, Z. Phys. C – Particles and Fields **48**(1990)159; Int. J. Mod. Phys. **A6**(1991)3081.
- [18] W. B. Schmidke, J. Wess and B. Zumino, Z. Phys. C – Particles and Fields **52**(1991)471.
- [19] O. Ogievetsky, W. B. Schmidke, J. Wess and B. Zumino, Lett. Math. Phys. **23**(1991)233.
- [20] O. Ogievetsky, W. B. Schmidke, J. Wess and B. Zumino, Commun. Math. Phys. **150**(1992)495.
- [21] J. Lukierski, H. Ruegg, A. Nowicki and V. N. Tolstoy, Phys. Lett. **264B**(1991)331.
- [22] J. Lukierski and A. Nowicki, Phys. Lett. **279B**(1992)299.
- [23] J. Lukierski, A. Nowicki and H. Ruegg, “Quantum Deformation of Poincaré and Conformal Algebras ”, Preprint BUHEP-91-21(1991).
- [24] T. Kobayashi and T. Uematsu, “q-Deformed Conformal and Poincaré Algebra on Quantum 4-spinors”, Preprint KUCP-52-revised(1993).
- [25] T. Kobayashi and T. Uematsu, Z. Phys. C – Particles and Fields **56**(1992)193.
- [26] P. van Nieuwenhuizen, Phys. Reports **68**(1981)189.
M. F. Sohnius, Phys. Reports **128**(1985)39.
- [27] E. Celeghini, R. Giacchetti, E. Sorace and M. Tarlini, J. Math. Phys. **31**(1990) 2548; **32**(1991) 1155; **32**(1991) 1159.